




# International Journal of Life Sciences Biotechnology and Pharma Research





Research Paper

## EFFECT OF POTENTIAL BIOCONTROL AGENTS AGAINST *SCLEROTIUM ROLFSII* CAUSING STEM ROT OF GROUNDNUT

K C Deepthi<sup>1\*</sup>

\*Corresponding Author: **K C Deepthi**, ✉ [Kc.deepthi@rediffmail.com](mailto:Kc.deepthi@rediffmail.com)

Total of 40 antagonists were isolated from groundnut rhizosphere and root endophytes of Regional agricultural research station of tirupati region. When tested *in vitro*, the potential biocontrol agents GRE-9 (100%) and GRB-16 (85%) were predominantly aggressive in inhibiting the mycelial growth of the pathogen in dual culture against *Sclerotium rolfsii*. Among the fungicides tested, Mancozeb was most compatible with all antagonists.

**Keywords:** Groundnut, Stem rot, *Sclerotium rolfsii*, Biocontrol agents

### INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is a major legume and oil seed crop in India, covering nearly half of the area under oilseeds. It is grown in over 100 countries with a total estimated area of 21.8 m. ha and with production of 28.5 MT. In India it is grown over an area of 4 lakhs/hectare, with an annual production of 5.5 million tonnes and yield was 1007 kg/hectare in the year 2009-10 (Economic survey 2010-11). *Sclerotium rolfsii* Sacc. is an important soil-borne pathogen and causes disease in numerous crops including peanut (Punja, 1988; Krupa and Dummergues, 1979). The loss of yield caused by pathogen infection generally is 25%, but sometimes it reaches 80-90% in some cases (Grichar and Bosweel, 1987). The disease causes damage on root and stem of plant. The pathogen produces

sclerotia which overwinter in soil and on plant debris and can survive in a long period causing disease in the following season (Punja, 1985). Thus, the control of the disease is very difficult. Different control methods have been investigated before including cultural practices (Gurkin and Jenkins, 1985; Punja *et al.*, 1986), chemical control (Bowen *et al.*, 1992; Minton *et al.*, 1993; Culbreath *et al.*, 1992), biological control (Elad *et al.*, 1984; Papavizas and Lewis, 1989; Latunde-Dada, 1993; and integrated control (Bicici *et al.*, 1994; Cilliers *et al.*, 2003). It is widely recognised that biological control of plant pathogens is the most important alternative for the future and can be exploited within the frame work of integrated disease management. The present study is mainly focussed to study and to isolate and identify potential antagonists of *S.rolfsii* and their

<sup>1</sup> Department of plant pathology, S.V. Agricultural College, Tirupati-517502.

compatibility with the commonly used fungicides.

## MATERIALS AND METHODS

### Isolation and Identification of Pathogen

The pathogen was isolated from stem rot infected groundnut plants showing typical symptoms of stem rot disease by tissue segment method (Rangaswami and Mahadevan, 1999).

The pathogen was identified based on mycelial and sclerotial characters based on standard mycological keys (Barnett and Hunter, 1972) and maintained on Potato dextrose agar (Dhingra and Sinclair, 1995) for further studies.

### Isolation of Root and Endophytes

For isolation of root endophytes and rhizosphere bacteria, 5 g of root were surface sterilized for 5 min. with 70.0 per cent ethanol and homogenized in 20 ml of sterilized phosphate buffer (20mM Na<sub>2</sub>HPO<sub>4</sub>+20mM NaH<sub>2</sub>PO<sub>4</sub>) using mortar and pestle. Appropriate dilutions (10<sup>-4</sup> for fungi and 10<sup>-6</sup> for bacteria) of these suspensions were plated on PDA and NA for the isolation of fungi and bacteria respectively. The plates were incubated for 72 hr at 28±2°C (Kishore *et al.*, 2005).

### Screening of Native Potential Biocontrol Agents by Dual Culture Technique

The antagonistic activity of microflora isolates against *S.rolfsii* was determined by dual culture technique (Johnson *et al.*, 1959). Mycelial discs measuring 6 mm diameter from four day old cultures of both fungal antagonist and the test pathogen were placed at equidistant on sterile Petri plate containing PDA medium. One day old cultures of bacteria were streaked on opposite side of the pathogen on PDA medium. The Petri plates were then incubated at 28 ± 2°C. Three replications were maintained in each treatment.

Suitable controls were kept without antagonist. Growth of antagonists, pathogen and zone of inhibition were measured after recording full growth in control plate. Percent inhibition of mycelial growth of test pathogen was calculated

### Efficacy and Compatibility of Native Potential Antagonists with Different Fungicides Under *In Vitro*

The selected native potential biocontrol agents were tested for their compatibility with the fungicides generally recommended for soil drenching viz., copper oxychloride (0.1%), mancozeb (0.2%), carbendazim (0.2%), thiophanate methyl (0.1%), Hexaconazole (0.1%) and Propiaconazole (0.1%). Fungal isolates were tested for their compatibility by poisoned food technique (Nene and Thapliyal, 1993) and spectrophotometric method (Kishore *et al.*, 2005) for bacterial isolates under *in vitro*.

### Statistical Analysis

Completely Randomized Design (CRD) was used for radial growth percent disease incidence, poisoned food technique and dual cultural technique (Gomez and Gomez, 1984)

## RESULTS AND DISCUSSION

A total of 40 antagonists were obtained from groundnut rhizosphere bacteria and root endophytes. 5 fungi and 40 bacterial isolates were obtained. The bacteria was designated as GRE 1 to GRE 20 (from Groundnut Root endophytes) and GRB 1 to GRB 20 (from Rhizosphere soil) similarly the fungi were named as GREF and GRBF and these were identified as *Trichoderma* spp. in which GRHF 75% from rhizosphere was found to be superior when compared to other isolates in inhibiting the growth of *S.rolfsii* in dual culture (Table 1 and Figure 1). Fungi in the genus

**Table 1: *In Vitro* Efficacy of Rhizosphere and Root Antagonistic by Trichoderma Isolates Against *S. rolfsii* in Dual Culture Technique**

Isolate	*Linear Growth of <i>S. rolfsii</i> (cm)	Percent of Inhibition of Mycelial Growth of <i>S. rolfsii</i>
GRAF-1	4	55.55 (48.18)
GRAF-2	4.5	50 (45.00)
GRAF-3	5	44.44 (41.80)
GRHF-4	2.2	75.55 (60.36)
GRHF-5	4.8	46.66 (43.08)
Control	9	0
CD (0.05)	-	7.99
S.Em±	-	2.59

Note: \* Mean of three replications.

**Figure 1: *In Vitro* Efficacy of Rhizosphere and Root Antagonistic *Trichoderma* Isolates Against *S. Rolfzii* in Dual Culture**



*Trichoderma* have been known since 1920s for the ability to act as biocontrol agents against plant pathogens (Harman, 2006). In case of 40 bacterial

isolates GRE 9 Completely inhibited (100%) followed by GRB 16 (85%) the growth of pathogen (Table 2).

**Table 2: In Vitro Evaluation of the Antagonistic Activity of Bacterial Root Endophytes and Rhizosphere Bacteria by *S. rolfsii* in Dual Culture Technique**

Isolate	*Linear Growth of <i>S. rolfsii</i> (cm)	Percent Inhibition of Mycelial Growth of <i>S. rolfsii</i>	Isolate	*Linear Growth of <i>S. rolfsii</i> (cm)	Percent Inhibition of Mycelial Growth of <i>S. rolfsii</i>
GRE-1	2.1	77.77 (61.86)	GRB-1	2.1	77.77 (61.86)
GRE-2	2.2	75.55(60.36)	GRB-2	4.3	52.22 (46.27)
GRE-3	2.0	76.66 (61.11)	GRB-3	3.8	57.77 (46.89)
GRE-4	3.7	58.88 (46.53)	GRB-4	3.0	65.55 (54.05)
GRE-5	3.1	66.66(54.73)	GRB-5	3.1	66.66 (54.73)
GRE-6	2.8	68.89 (56.09)	GRB-6	2.6	71.11(57.48)
GRE-7	2.43	72.96 (58.83)	GRB-7	3.5	61.11 (51.26)
GRE-8	2	77.78 (61.88)	GRB-8	2	77.78 (61.88)
<b>GRE-9</b>	<b>0.0</b>	<b>100 (90.00)</b>	GRB-9	3.2	64.44 (53.39)
GRE-10	2.83	68.52 (55.87)	GRB-10	2.83	68.52 (55.87)
GRE-11	3.96	55.93 (48.41)	GRB-11	3.9	55.93 (48.41)
GRE-12	3.13	65.19 (53.84)	GRB-12	3.1	65.19 (53.84)
GRE-13	2.5	72.22 (58.19)	GRB-13	2.5	72.22 (58.19)
GRE-14	3.4	62.00 (51.94)	GRB-14	3.4	62.00 (51.94)
GRE-15	3.6	60 (50.76)	GRB-15	3.6	60 (50.76)
GRE-16	2.2	75.55 (60.36)	<b>GRB-16</b>	<b>1.3</b>	<b>85.55 (67.65)</b>
GRE-17	3.0	66.66 (54.73)	GRB-17	3.0	66.66 (54.73)
GRE-18	4.0	57.23 (49.34)	GRB-18	2.03	77.41 (61.66)
GRE-19	2.43	72.96 (58.83)	GRB-19	3.3	63.33 (52.73)
GRE-20	1.4	84.44 (66.76)	GRB-20	1.8	80.00 (63.43)
Control	9	0	Control	9	0
CD(0.05)	-	10.21	CD(0.05)	-	7.97
S.Em±	-	3.46	S.Em±	-	2.70

Note: \* Mean of three replications.

The use of biological control methods to reduce disease incidence caused by plant pathogens is continually being developed and is

being used in variety of crops. By including biocontrol agents for controlling *S.rolfsii* as a part of integrated disease management will help in

managing disease as well as delaying the development of tolerance/resistance to fungicides. The bacterial isolate GRE 9 (Table 3) was more compatible with mancozeb followed by carbendazim, Copper oxychloride. Similar observations were made by Vidhyasekharan and Muthamilan (1995) and reported that carbendazim was not inhibitory to *P. fluorescens*. The fungal isolate GRHF4 (Table 4 and Figure 2) was more

compatible with Mancozeb followed by copper oxychloride. Similar results were obtained by Vijayaraghavan and Abraham (2004) observed that mancozeb was compatible with *T.viride*. Among all the fungicides tested Mancozeb was found to be more compatible. The present study revealed that the pathogen *S.rolfsii* developed tolerance against mancozeb when compare to other fungicides.

**Table 3: In Vitro Evaluation of the Compatibility of the Potential Antagonistic Bacterial Isolate Gre-9 With Different Fungicides**


Fungicide	Concentration	*Growth of Bacterial Isolate (OD at 600nm)
Mancozeb	0.2%	1.45
Carbendazim	0.2%	0.92
Copper oxychloride	0.1%	0.86
Thiophanate methyl	0.1%	0.67
Hexaconazole	0.1%	0.78
Propiaconazole	0.1%	0.80
Control	-	1.56
CD (0.05)	-	0.27
SEm±	-	0.096

Note: \* Mean of three replications.

**Table 4: In Vitro Evaluation of Compatibility of Trichoderma Isolate GRHF-4 With Different Fungicides in Poisoned Food Technique**

Fungicide	Concentration	*Growth of Trichoderma (mm)	Per cent inhibition over control
Mancozeb	0.2%	4.7	48.01
Copper oxychloride	0.2%	2.1	77
Carbendazim	0.1%	0	100
Thiophanate methyl	0.1%	0	100
Propiaconazole	0.1%	0	100
Hexaconazole	0.1%	0.1	99
Control		9.0	-
CD(0.05)		-	5.52
SEm±		-	1.82

**Figure 2: *In Vitro* Evaluation of Compatibility of Tricoderma Isolate GRBF-4 With Different Fungicides in Poisoned Food Technique**



## CONCLUSION

The results revealed that GRE -9 has showed 100% inhibition against the pathogen (*S.rolfsii*). The bacterial isolate GRE 9 was more compatible with mancozeb under poison food technique. In future the work will be carried on integrated disease management to reduce inoculum thresh hold levels and to suppress the disease for the purpose of higher yields.

## ACKNOWLEDGMENT

One of the authors expresses sense of gratitude to Prof. N.P. Eswara Reddy, Department of plant pathology, S.V. Agricultural college, Tirupati for providing laboratory facilities and kind support during the course of these investigations.


## REFERENCES

1. Barnett H L and Barry B Hunter (1972), *Illustrated genera of imperfect fungi*, Burgess publishing company, Minnesota.
2. Bicici M, Cinar O and Erkilic A (1994), "The control of stem rot caused by *Sclerotium rolfsii* Sacc on peanut by cultural, chemical, physical and biological methods", *Tr J. Agric. For.*, Vol. 18, pp. 423-435.
3. Bowen K L, Hagan A K and Weeks R (1992), "Seven years of *Sclerotium rolfsii* in peanut fields: yield losses and means of minimization", *Plant Dis.*, Vol. 76, pp. 982-985.
4. Cilliers A J, Pretorius J A and Van Wyk P S

- (2003), "Integrated control of *Sclerotium rolfsii* on groundnut in South Africa", *J. Phytopathol.*, Vol. 151, pp. 249-258.
5. Culbreath A K, Minton N A, Brenneman T B and Mullinix B G (1992), "Response of florunner and southern runner peanut cultivars to chemical management of late leaf spot, southern stem rot and nematodes", *Plant Dis.*, Vol. 76, pp. 1199-1203.
  6. Dhingra O D and Sinclair J B (1995), *Basic Plant Pathology Methods*, CRC Press London.
  7. Elad Y, Barak R and Chet I (1984), "Parasitism of sclerotia of *Sclerotium rolfsii* by *Trichoderma harzianum*", *Soil Biol. Biochem.*, Vol. 16, pp. 381-386
  8. Gomez K A and Gomez A A (1984), *Statistical procedures for agricultural research*, Second Edition, John Wiley and Sons, New York.
  9. Grichar V J and Bosweel T E (1987), "Comparison of lorsban and tilt with terrachlor for control of southern blight on peanut", *The Texas Agriculture Experiment Station*, PR-4534.
  10. Gurkin R S and Jenkins S F (1985), "Influence of cultural practices, fungicides and inoculum placement on southern blight and Rhizoctonia crown rot of carrot", *Plant Dis.*, Vol. 69, pp. 477-481.
  11. Harman G E (2006), "Over view of mechanisms and uses of *Trichoderma* spp.", *Phytopathology*, Vol. 96, pp. 190-194.
  12. Kishore G K Pande S and Podile A R (2005), "Biological control of collar rot disease with broad-spectrum antifungal bacteria with groundnut", *Canadian Journal of Microbiology*, Vol. 51, pp. 123-132.
  13. Latunde-Dada A O (1993), "Biological control of southern blight disease of tomato caused by *Sclerotium rolfsii* with simplified mycelia formulations of *Trichoderma koningii*", *Plant Pathol.*, Vol. 42, pp. 522-529.
  14. Minton N A, Brenneman T B, Bondari K and Harrison G W (1993), "Activity of fosthiazate against *Meloidogyn aenaria*, *Frankliniella* sp. And *Sclerotium rolfsii* in peanut", *Peanut Sci.*, Vol. 20, pp. 66-70.
  15. Nene Y L and Thapliyal P N (1993), *Fungicides in plant disease control*, 3rd Edition, Oxford and IBH publishing company, New Delhi.
  16. Papavizas D C and Lewis J A (1989), "Effect of Gliocladium and Trichoderma on damping-off and blight of snapbean caused by *Sclerotium rolfsii* in the greenhouse", *Plant Pathol.*, Vol. 38, pp. 277-286.
  17. Punja Z K (1985), "The biology, ecology and control of *Sclerotium rolfsii*", *Ann Rev Phytopathol.*, Vol. 23, pp. 97-127.
  18. Punja Z K (1988), "*Sclerotium (Athelia) rolfsii* A Pathogen of Many Plant Species", in Sidhu G S (Ed.), *Advances in Plant Pathology*, Vol. 6 Genetics of Plant Pathogenic Fungi, Academic Press, London, pp. 523-534.
  19. Punja Z K, Carter J D, Campell G M and Rossell E L (1986), "Effects of calcium and nitrogen fertilizers, fungicides and tillage practices in incidence of *Sclerotium rolfsii* on processing carrots", *Plant Dis.*, Vol. 70, pp. 819-824.



20. Rangaswami G and Mahadevan A (1999), *Diseases of crop plants in India*, 4th Edition, Prentice Hall of India Pvt. Ltd. New Delhi, p. 6079.
21. Vidyasekaran P and Muthamilan (1995), "Development of formulations of *Pseudomonas fluorescens* for control of chickpea wilt", *Plant Diseases*, Vol. 79, pp. 782-786.
22. Vijayaraghavan R and Abraham K (2004), "Compatibility of biocontrol agents with pesticides and fertilizers used in black pepper gardens", *Journal of Mycology and Plant Pathology*, Vol. 34, pp. 506-510.



**International Journal of Life Sciences Biotechnology and Pharma Research**

**Hyderabad, INDIA. Ph: +91-09441351700, 09059645577**

**E-mail: editorijlbpr@gmail.com or editor@ijlbpr.com**

**Website: www.ijlbpr.com**

