



# International Journal of Life Sciences Biotechnology and Pharma Research





## Research Paper

# MOLECULAR CHARACTERIZATION AND METHYLOTROPHIC ACTIVITIES OF *PSEUDOMONAS* SPP. FROM LONAR LAKE

Tambekar D H<sup>1\*</sup> and A L Pawar<sup>1</sup>\*Corresponding Author: Tambekar D H, ✉ [diliptambekar@rediffmail.com](mailto:diliptambekar@rediffmail.com)

Lonar Lake is a saline ecosystem formed entirely on basalt rock. It is a completely closed system a favors the growth of alkalophiles and halophiles. The *Pseudomonas* spp play a major role for conserving environment by utilizing toxic C<sub>1</sub> carbon compound and reduces the pollution produced by methanol, green houses gases etc. In present study, methylotrophic bacterial isolates were isolated from water and sediment of Lonar Lake using minimal salt medium containing 2% (v/v) methanol as sole carbon and energy source. Total six bacterial strains were isolated and characterized morphologically, biochemically and identified as *Pseudomonas aeruginosa* by 16S rRNA sequencing. These *Pseudomonas* strains were further screened for its ability to utilize methanol by Spectrophotometric method. Results showed that the isolates ALP4 and ALP6 of *Pseudomonas* strains are found to be highly efficient methanol utilizer and could be effectively use for bioremediation by reducing methanol at polluted sites and also helpful in reducing C<sub>1</sub> compound, and global warming gases from environment.

**Keywords:** Lonar Lake, Methylotrophs, Bioremediation, Methanol and *Pseudomonas*

## INTRODUCTION

The Lonar Lake is a saline and hyperalkaline ecosystem formed entirely on basalt rock by meteor impact. The crater is located in India ~ 550 km east of Mumbai and Arabian Sea is 150 m deep and 1830 m across (Fredrikson, 1973) with raised rim up to the 100m in width and 20m to 30 m high and alkalinity of the Lake water is attributed to the high content of sodium carbonate (Thakker and Ranade, 2002). The Lonar Lake is unique in the world for its alkalinity (pH 10) and salinity (NaCl 0.9%) of the water but it was seen

that chlorides and salinity of the Lake water is decreasing day by day (Tambekar *et al.*, 2010). Microbiological studies using culture-dependent and culture-independent strategies have identified and characterized both bacterial (Kanekar *et al.*, 1999, 2002; Nilegaonkar *et al.*, 2002; Wani *et al.*, 2006; Surakasi *et al.*, 2007) and archaeal (Thakker and Ranade, 2002; Surakasi, 2007; Surakasi *et al.*, 2007) communities in the Lonar Lake water and sediment.

Methanotrophs are a unique group of methylotrophic bacteria, which utilize one-carbon

<sup>1</sup> Sant Gadge Baba Amravati University, Amravati - 444602 (MS)

(C<sub>1</sub>) compounds such as methane, methanol and methylamine which constitute an important component of microbe-driven food web chains in many ecosystems. Methylophilic bacteria, are phylogenetically distributed across diverse phyla such as *Gammaproteobacteria* (type I methanotrophs), *Alphaproteobacteria* (type II methanotrophs) (Trotsenko and Murrell, 2008), filamentous methane oxidizers (Stoecker *et al.*, 2006; Vigliotta *et al.*, 2007) and *Verrucomicrobia* (Dunfield *et al.*, 2007; Pol *et al.*, 2007; Islam *et al.*, 2008) and contribute significantly in biogeochemical cycling of carbon by facilitating the incorporation of C<sub>1</sub> compound-derived carbon into biomass (Anthony, 1992; Chistoserdova *et al.*, 2009). [HYPERLINK "http://www.nature.com/ismej/journal/v4/n11/full/ismej20107"](http://www.nature.com/ismej/journal/v4/n11/full/ismej20107)[HYPERLINK "http://www.nature.com/ismej/journal/v4/n11/full/ismej2010"](http://www.nature.com/ismej/journal/v4/n11/full/ismej2010)The global cycling of methane and related C<sub>1</sub> compounds further affects the important environmental phenomena related to climate change.

The toxicity of methanol has been widely documented and their disastrous effect towards human and environment is greatly concerned. Currently utilization of C<sub>1</sub> compound received great attention from many people from industries and researcher due to their toxicity. The present study aimed to isolate and characterize methylophilic bacterial spp from Lonar Lake which can remediate methanol by which there may reduction in C<sub>1</sub> compound pollution in the environment. The phylogenetic analysis of these methylophilic bacterial isolates could be very much helpful in future research, related to microbial diversity and reduction in pollution level of global gases from environment.

## MATERIALS AND METHODS

### Isolation of Bacterial Strain by Enrichment Method

The sediment and water samples were collected from different sites of Lonar Lake (Buldhana District in Maharashtra, India). The medium containing ingredient (g/l): NaNO<sub>3</sub> 2.5g, KCl 0.1g, KH<sub>2</sub>PO<sub>4</sub> 3g, K<sub>2</sub>HPO<sub>4</sub> 7g, CaCl<sub>2</sub> 0.01g, MgSO<sub>4</sub> 0.5g, FeSO<sub>4</sub> 0.116g, H<sub>3</sub>BO<sub>3</sub> 0.232g, CuSO<sub>4</sub> 0.41g, MnSO<sub>4</sub> 0.008g, (NH<sub>4</sub>)<sub>6</sub> Mo<sub>7</sub>O<sub>24</sub>, 0.008g, and ZnSO<sub>4</sub> 0.174g, 20 ml methanol, is used for isolation of methylophilic bacteria (Haddad *et al.*, 2009) The medium was then incubated at 37°C on a rotary shaking incubator at 100 rpm for 3 days. The sub-culturing was made for 5 times for enrichment and isolation of pure culture was done on solid nutrient agar plate. Well isolated colonies were selected and stored at 4°C as stock culture (Tambekar *et al.*, 2010).

### Identification of Isolates

The isolates were characterized by morpho-physiological and cultural characteristics. The biochemical tests were performed by rapid detection kit (Hi-media, KB001 and KB009). The kit contents the tests namely, indol, methyl red, voges prauskar, citrate, lactose, xylose, maltose, fructose, dextrose, galactose, raffinose, trehalose, melibiose, sucrose, L-arabinose, mannose, inulin, sodium gluconate, glycerol, salicin, glucosamine, dulcitol, inositol, sorbitol, Rhamnose, cellobiose, melizitiose, α-methyl mannose, xylitol, ONPG, esculin, arabinose, citrate, malonate, sorbase, nitrate reduction, urea hydrolysis and starch hydrolysis. The molecular identifications were performed by 16S rRNA sequencing at NCCS, Pune.

### Methanol Utilization Studies

For estimation of methanol, the bacterial isolates

were grown in the nutrient broth and incubated overnight at 37 °C. This 24 h old culture was inoculated in mineral salt medium containing 2% (v/v) methanol as a sole source of carbon and energy. Preliminary studies were carried out bacterial inoculums in medium containing 2% (v/v) methanol. The methanol utilization was determined by analyzing residual methanol at 480 nm in the medium after the each interval 24 h up from 0 h to 96 h by using UV- visible spectrophotometer (Zhan *et al.*, 2010).

## RESULTS AND DISCUSSION

Lonar Lake represents an extreme environment with high pH and moderate salinity. It is the only known depression in the region and hence may serve as a drain for excess runoff from anthropogenically influenced surrounding areas. However, the contribution of such natural or anthropogenic factors towards elevated phosphate and nitrate levels in the lake sediments warrants further investigation. Lonar Lake water is green throughout the year because of dense cyanobacterial bloom dominated by *Arthrospira* (Surakasi *et al.*, 2007). Sediment and water samples were chosen as the source of bacterial isolation and these were enriched in minimal medium using 2% (v/v) methanol as sole source of carbon and energy for one month by repeated subculture after every 96 h.

Those bacterial isolates able to grow on medium containing 2% (v/v) methanol as carbon source were identified as methylotrophs and subsequently isolated in pure culture. The experimental outcome of morphological and biochemical characterization proved that all isolates are gram negative and identified as *Pseudomonas* (Table 1) and represented as ALP1, ALP 2, ALP 4, ALP 5, ALP 6 and ALP 9.

These isolated were analysed for 16s rRNA and identified as *Pseudomonas aeruginosa*. The *Pseudomonas aeruginosa* have best potential to utilize methanol. These isolates were able to grow at temperature up to 42°C.

In this investigation, a new method for the direct determination of methanol using sodium nitroprusside (SNP) is used (Zhan *et al.*, 2010). It has been reported that SNP can react with nucleophilic agent such as primary and secondary aliphatic amine however; no studies in the literature to date have been reported on the reaction of SNP and alcohol. This experiment results showed that SNP can react with methanol to form colored product. Absorbance of product is linear with certain extent of the concentration of methanol.

These morpho-biochemically characterized bacteria were identified by 16S rRNA sequencing and phylogenetic tree was constructed. The result of the phylogeny showed that methylotrophic strains isolated from Lonar Lake were related to phylum Proteobacteria. According to 16S rRNA gene sequences, all these isolates showed a high level of similarity with the genus *Pseudomonas* (Figure 1).

Sediment is the most potential source to isolate *Pseudomonas*. Antony *et al.*, (2010) identified methylmicrobium, methylphaga and *Bacillus* spp as predominant methanotrophs from sediments of Lonar Lake. The utilization performances of selected strain were examined by spectrophotometric method. The experiment designed to find out the percent utilization and rate of utilization after 24h, 48h, 72h and 96h. In these studies, methanol estimated for 96h by spectroscopic method at each 24h time interval. Percent utilization and rate of degradation of

**Table 1: Morphological, Cultural and Biochemical Characteristic of Bacteria Isolated From Lonar Lake**


Culture Code →	ALP 1	ALP 2	ALP 4	ALP 5	ALP 6	ALP 9	Culture Code →	ALP 1	ALP 2	ALP 4	ALP 5	ALP 6	ALP 9
Bacterial isolates on the basis of 16S rRNA sequencing →	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas</i> spp.	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>	Bacterial isolates on the basis of 16S rRNA sequencing →	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas</i> spp.	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i>
Source	Sed	W	W	Sed	Sed	Sed	Dulcitol	-	-	-	-	-	-
Colony Colour	Gr	Gr	Gr	Gr	Wh	Gr	Glycerol	-	-	-	-	-	-
Colony shape	Ir	Ir	Ir	Ir	Ir	Ir	Salicin	-	-	-	-	-	-
Gram Reaction	-	-	-	-	-	-	Glucosamine	-	-	-	-	-	-
Shape of bacteria	SR	SR	SR	SR	SR	SR	Sodium gluconate	-	-	-	-	-	-
Arrangement	S	S	S	S	S	S	Inositol	-	-	-	-	-	-
Endospore	-	-	-	-	-	-	Sorbitol	-	-	-	-	-	-
Motility	+	+	+	+	+	+	Mannitol	-	-	-	-	-	-
Catalase	-	-	-	-	-	-	Adonitol	-	-	-	-	-	-
Oxidase	+	+	+	+	+	+	$\alpha$ -Methyl d-glucoside	-	-	-	-	-	-
Indol	-	-	-	-	-	-	Ribose	-	-	-	-	-	-
Methyl red	-	-	-	-	-	-	Rhamnose	-	-	-	-	-	-
Voges Praskaur	-	-	-	-	-	-	Cellibiose	-	-	-	-	-	-
Citrate	+	+	+	+	+	+	Melizitose	-	-	-	-	-	-
Lactose	-	-	-	-	-	-	$\alpha$ -Methyl d-mannose	-	-	-	-	-	-
Xylose	+	+	+	+	+	-	Xylitol	+	+	+	+	+	+
Maltose	-	-	-	-	-	-	ONPG	-	-	-	-	-	-
Fructose	-	-	-	-	-	-	Esculin	-	-	-	-	-	-
Dextrose	+	+	+	+	+	+	D-arabinose	-	-	-	-	-	-
Galactose	+	+	+	+	+	-	Malonate	+	+	+	+	+	+
Raffinose	-	-	-	-	-	-	Sorbose	-	-	-	-	-	-
Trehalose	-	-	-	-	-	-	Nitrate Reduction	-	-	-	-	+	-
Melibiose	-	-	-	-	-	-	Urease	-	-	-	-	-	-
Sucrose	-	-	-	-	-	-	Starch Hydrolysis	-	-	-	-	-	-
L-arabinose	-	-	-	-	-	-	Growth at 6.5% NaCl	+	+	+	+	+	+
Mannose	+	+	+	+	+	+	Growth at 4°C	-	-	-	-	-	-
Inulin	-	-	-	-	-	-	Growth at 42°C	+	+	+	+	+	+

Note: Sed-Sediment, W-Water, Gr-Green, Wh-white, I-Irregular, SR-Short rod, S-Single.

methanol for all these isolates was found almost similar in range 16% to 74% and 0.033 mg/l to 0.045mg/l respectively with time interval 24h, 48h,


72h, 96h. Experiment also showed that out of these isolates *Pseudomonas* ALP1 utilize 72% of methanol, ALP 2 utilize 70% of methanol, ALP

**Figure 1: Phylogenetic Tree of *Pseudomonas* Species Isolated from Lonar Lake**




4 utilize 72 % of methanol, ALP 5 utilize 68 % of methanol, ALP 6 utilize 74% of methanol and ALP 9 utilized 70% of methanol after 96h and rate of utilization of methanol was about 0.04 for all isolates of *Pseudomonas* spp (Figure 2). The rate of utilization of methanol was also almost same for all bacterial isolates (Figure 3). Tambekar *et al.*, (2011) reported that *Pseudomonas aeruginosa* is new species and not previously recorded bacterial species from Lonar Lake to utilize methanol as carbon source. Previous records indicated that *Pseudomonas aeruginosa* found in soil and water, and involved in

**Figure 2: Percent Utilization of Methanol by *Pseudomonas* Species**



**Figure 3: Rate of Degradation of Methanol by *Pseudomonas* Species**



bioremediation. The findings of this study provide a window into the diversity of bacterial community members which are methane degrading from the Lonar Lake. These isolated bacterial species may be used to combat industrial pollution of methanol or to control global warming which may found better choice for studies like methane, methanol or toxic chemical degradation to combat Global warming. Till date several works are in progress to isolates efficient microbial strain that have ability to utilize methanol. We report here a new *Pseudomonas aeruginosa* (ALP4 and ALP6) was found to be a potential strain to utilize methanol as sole source of carbon and energy. This work has provided a useful guideline in evaluating potential methanol utilizer isolated from Lonar Lake.

## CONCLUSION


The unexplored site of alkaline Lonar Lake contains many methanogenic and methylotrophic genera which might be helpful for the remediation of pollution environment. In present study isolation strategy for methylotrophic bacteria was used and potential methanol utilizing bacterial isolates of *Pseudomonas* species were isolated which can be helpful for remediation of site with pollution of C1 compounds and provides a new unexplored site for researcher.

## REFERENCES

1. Anthony C (1992), "The Biochemistry of Methylotrophs", Academic Press, New York.
2. Antony C V, Kumarsan D, Ferrando L, Boden H, Moussard H, Scavino AF, Shouche Y S and Murrel J C (2010), "Active methylotrophs in the sediments of Lonar Lake, a Saline and Alkaline Ecosystem formed by Meteor Impact", *The ISME J.*, Vol. 4, pp.1470-1480.
3. Chistoserdova L, Kalyuzhnaya M G and Lidstrom M E (2009), "The Expanding World of Methylotrophic Metabolism", *Ann. Rev. Microbiol.*, Vol. 63, pp. 477-499.
4. Dunfield P F, Yuryev A, Senin P, Smirnova A V, Stott M B and Hou S (2007), "Methane oxidation by an extremely acidophilic bacterium of the phylum *Verrucomicrobia*", *Nat.*, Vol. 450, pp. 879-882.
5. Fredriksson K, Dube A, Milton DJ and Balasundaram M S (1973), "Lonar Lake, India: An Impact Crater in Basalt", *Science.*, Vol. 180, pp. 862-864.
6. Haddad N I A, Wang J and Bozhong M (2009), "Identification of biosurfactant producing strain: *Bacillus subtilis* HOB2", *Protein and Peptide Lett.*, Vol.16, pp. 7-13.
7. Islam T, Jensen S, Reigstad L J, Larsen O and Birkeland N K (2008), "Methane oxidation at 55°C and pH 2 by a thermoacidophilic bacterium belonging to the *Verrucomicrobia* phylum", *Proc Natl Acad Sci USA.*, Vol. 105, pp. 300-304.
8. Kanekar P P, Nilegaonkar S S, Sarnaik SS and Kelkar A S (2002), "Optimization of protease activity of alkaliphilic bacteria isolated from an alkaline lake in India", *Bioresour Technol.*, Vol. 85, pp. 87-93.
9. Kanekar P P, Sarnaik S S and Kelkar A S (1999), "Bioremediation of Phenol by Alkaliphilic Bacteria Isolated from Alkaline Lake of Lonar, India", *J. Appl. Microbiol.*, Vol. 85: pp.128-133.
10. Nilegaonkar S S, Kanekar P P, Sarnaik S S and Kelkar A S (2002), "Production, isolation and characterization of extracellular protease of an alkaliphilic strain of

- Arthrobacter ramosus*, MCM B-351 isolated from the alkaline lake of Lonar, India”, *World J. Microbiol. Biot.*, Vol.18, pp. 785-789.
11. Pol A, Heijmans K, Harhangi H R, Tedesco D, Jetten M S and Op den Camp H J (2007), “Methanotrophy below pH 1 by a new *Verrucomicrobia* species”, *Nature*, Vol. 450, pp. 874-878.
  12. Stoecker K, Bendinger B, Schoning B, Nielsen P H, Nielsen J L and Baranyi C (2006), “Cohn2s *Crenothrix* is a filamentous methane oxidizer with an unusual methane monooxygenase”, *Proc. Natl. Acad. Sci. USA.*, Vol. 103, pp. 2363-2367.
  13. Surakasi V P (2007), “Diversity of Methanogenic Archaea of Lonar Lake”, Ph. D. thesis University of Pune: Pune, India.
  14. Surakasi V P, Wani A A, Shouche Y S and Ranade D R (2007), “Phylogenetic Analysis of Methanogenic Enrichment Cultures Obtained from Lonar Lake in India: isolation of *Methanocalculus* sp. and *Methanoculleus* sp.”, *Microb. Ecol.*, Vol. 54, pp. 697-704.
  15. Tambekar D H, Patil R V and Pawar A L (2011), “Studies on methanotrophs from Lonar Lake”, *J. research in Biol.*, Vol. 3, pp. 230-236.
  16. Tambekar D H, Pawar A L and Dudhane MN (2010), “Lonar Lake Water: Past and Present”, *Nat. Environ. Poll Technol.*, Vol. 9, No. 2, pp. 217-221.
  17. Thakker C D and Ranade D R (2002), “Alkalophilic *Methanosarcina* Isolated from Lonar Lake”, *Curr. Sci.*, Vol. 82, pp. 455-458.
  18. Trotsenko YA and Murrell JC (2008), “Metabolic Aspects of Aerobic Obligate Methanotrophy”, *Adv. Appl. Microbiol.*, Vol. 63: pp.183-229.
  19. Vigliotta G, Nutricati E, Carata E, Tredici S M, De S M and Pontieri P (2007), “*Clonothrix fusca* Roze 1896, a filamentous, Sheathed, Methanotrophic Gamma-proteobacterium”, *Appl. Environ. Microbiol.*, Vol. 73, pp. 3556-3565.
  20. Wani A A, Surakasi V P, Siddharth J, Raghavan R G, Patole M S and Ranade D (2006), “Molecular Analyses of Microbial Diversity Associated with the Lonar Soda Lake in India: An impact crater in a basalt area”, *Res. Microbiol.*, Vol. 157, pp. 928-937.
  21. Zhan Y Y, Zhang Y, Li Q M and Du X Z (2010), “A Novel Visible Spectrophotometer Method for Determination of Methanol Using Sodium Nitroprusside as Spectroscopic Probe”, *J. Chinese Society*, Vol. 57, pp. 230-235.





**International Journal of Life Sciences Biotechnology and Pharma Research**

**Hyderabad, INDIA. Ph: +91-09441351700, 09059645577**

**E-mail: editorijlbpr@gmail.com or editor@ijlbpr.com**

**Website: www.ijlbpr.com**

