



# International Journal of Life Sciences Biotechnology and Pharma Research





Review Article

## SEED DETERIORATION: A REVIEW

Jyoti<sup>1</sup> and C P Malik<sup>1\*</sup>

\*Corresponding Author: **C P Malik**, ✉ [cpm\\_malik@yahoo.com](mailto:cpm_malik@yahoo.com)

Seed deterioration is an undesirable and detrimental attribute of agriculture. This process is a separate event from seed development and germination. Losses in seed quality occur during field weathering, harvesting and storage. Deterioration caused by field weathering is directly related to seed exposure to adverse conditions. Seeds are highly susceptible to damage and mechanical injury during post-harvest handling. Several environmental factors contribute to seed deterioration and these conditions make very difficult to maintain viability during storage. Seed quality is depends upon initial seed quality, temperature, moisture content and mycoflora. Rapid deterioration occurs due to these environmental conditions make very difficult to maintain its viability during storage. However, the seed quality and viability during storage depend upon the initial quality of seed and the manner in which it is stored. Seed deterioration is associated with various cellular, metabolic and chemical alterations including lipid peroxidation, membrane disruption, DNA damage, impairment of RNA and protein synthesis and cause several detrimental effects on seed.

**Keywords:** Deterioration, Field weathering, Lipid peroxidation, Membrane disruption

### INTRODUCTION

Seed deterioration is loss of seed quality, viability and vigor due to effect of adverse environmental factors (Kapoor *et al.*, 2010). Deteriorative changes enhance when seed exposure to external challenges increases and decrease the ability of the seed to survive. It is an undesirable attribute of agriculture. Annual losses due to deterioration can be as much as 25% of the harvested crop. It is one of the basic reasons for low productivity (Shelar *et al.*, 2008). The process has been described as cumulative, irreversible,

degenerative and inexorable process (Kapoor *et al.*, 2011). As seed deterioration increases, seed performance is progressively decreases. The physiology of seed deterioration is a distinguish consequence from seed development and germination.

Losses in seed quality occur during field weathering, harvesting and storage. Several factors contribute to the susceptibility for seed deterioration. The basic causes are temperature, relative humidity, seed moisture content and by invasion of and damage to tissues by

<sup>1</sup> School of Life Sciences, Jaipur National University, Jaipur, Rajasthan, India.

microorganisms, insects. The rate of deterioration fluctuates critically from one species to another and also among varieties of the same species (Jatoi *et al.*, 2001). Deterioration is evident as a reduction in percentage germination, produce weak seedlings, loss of vigor, become less viable and ultimately seed death (Maity *et al.*, 2000; Tilebeni and Golpayegani, 2011). The percentage emergence of deteriorated seeds is less than healthy seeds. Hence, deteriorated seed produces uneven stands, spotty fields, and fewer plants per hectare than healthy seed (Biabani *et al.*, 2011). Plants that have originated from deteriorate seed can also reduce growth rate (Kapoor *et al.*, 2010).

## WHAT IS DETERIORATION?

Seed deterioration can be defined as “deteriorative alterations occurring with time that increase the seed’s exposure to external challenges and decrease the ability of the seed to survive”. Seed deterioration causes loss of seed quality with time. It is a natural process which involves cytological, physiological, biochemical and physical changes in seeds. These changes reduce viability and ultimately cause death of the seed.

## TYPES OF DETERIORATION

Deterioration is evident as a decrease in percentage germination, while those seeds that germinate produce weak seedlings. Losses in seed quality occur during field weathering, harvesting and storage (Farhadi *et al.*, 2012). Harvesting time of several crops depends on its maturity time and on physiological maturity. Harvesting stage influences the quality of seed, germination, vigor, viability and also storability (Khatun *et al.*, 2009). Physiological maturity

attainment is a genotypic character which is influenced by several environmental factors (Mahesha *et al.*, 2001a).

Deterioration of seed in the field before harvest (field weathering) begins when the seed reaches physiological maturity and it extends till the seeds are harvested. The moisture content of physiological matured seed is approximately 50-55%. Because of the high moisture content, the seed cannot be harvested commercially and must remain in storage on the plant through a desiccation period till moisture levels are adequately low to permit mechanical harvest without causing undue damage to the seed. This desiccation period vary from a few days to several weeks before the seed attains a harvestable moisture level, about 14%. During this post-maturation, pre-harvest period weather conditions have a great influence on the quality of the harvested seed. Field conditions are rarely favorable for such storage. Seed quality is influenced by numerous factors that occur in the field before harvesting and during harvesting, drying, processing and storage. The losses are worsened if seeds are stored at high temperatures and high relative humidity conditions (Mosavi Nik *et al.*, 2011).

### Field Weathering

The deterioration of seed quality, vigor and viability, due to high relative humidity and high temperature during the post-maturation and pre-harvest period is referred to as field weathering (Bhatia *et al.*, 2010). Weathering occurs in the period between the attainment of physiological maturity till harvesting in the field. Deterioration caused by weathering is directly related to seed exposure to adverse conditions, so that the physiological

quality is depending on the environmental conditions preceding harvesting (Pádua *et al.*, 2009).

Exposure to hot and humid conditions, rainfall, photoperiod after ripening are pre-harvest factors, cause seed quality loss following physiological maturity. Among all these factors, influence of moisture on seeds during ripening appears to exert the major influence on predisposition to weathering. Adverse environmental conditions during seed filling and maturation result in forced seed maturation, which is associated with low yields, leading to a significant decrease in quality and an extensive reduction in the crop productivity (França-Neto *et al.*, 2005; Pádua *et al.*, 2009).

After physiological maturity if the seeds are retained on mother plant seeds will deteriorate, physiological changes in seed may lead to formation of rigid seeds or off color seeds in pulse crops (Khatun *et al.*, 2009). Harvest delays beyond optimum maturity extend field exposure and intensify seed deterioration. Weathering not only lowers seed germination, but also increases susceptibility to mechanical damage and disease infection. Timely harvesting avoids prolonged exposure to moisture, and is the best means of avoiding weathering.

### Harvest and Post-harvest Deterioration

Seed quality is highly affected by harvesting and handling methods. Harvest and post-harvest deterioration comprises threshing, processing machinery, seed collection, handling, transporting and drying. Mechanical damage is one of the major causes of seed deterioration during storage. Very dry seeds are prone to mechanical damage and injuries. Such damage may result in physical damage or fracturing of essential seed

parts; broken seed coats permit early entry and easy access for microflora, make the seed vulnerable to fungal attack and reduce storage potential (Shelar, 2008). In its severest form, physical seed damage is exhibited by splitting of the cotyledon, shattered and broken seeds. Large seeded varieties are more sensitive to mechanical damage than small seeds.

### Storage

Storability of seeds is mainly a genetically regulated character and is influenced by quality of the seed at the time of storage, pre-storage history of seed (environmental factors during pre- and post-harvest stages), moisture content of seed or ambient relative humidity, temperature of storage environment, duration of storage and biotic agents (Shelar *et al.*, 2008; Balešević-Tubić *et al.*, 2005; Khatun *et al.*, 2009; Biabani *et al.*, 2011). Damage of seed during storage is inevitable (Balešević-Tubić *et al.*, 2005).

These environmental conditions are very difficult to maintain during storage. The seed storage environment highly influences the period of seed survival. After planting of deteriorate seeds, seedling emergence may be poor and transmission of pathogens to the new crop may occur. Lower temperature and humidity result in delayed seed deteriorative process and thereby leads to prolonged viability period (Mohammadi *et al.*, 2011).

## MECHANISMS OF SEED DETERIORATION

Once seed deterioration has happened, this catabolic process cannot be reversed. It is a sequence of events beginning with a chain of biochemical events, predominantly membrane

damage and impairment of biosynthetic reactions, and then the resulting losses of various seed performance attributes, starting with reduced germination rate, reduced field emergence, increased numbers of abnormal seedlings and finally seed death. Viability loss results in irreversible chemical and structural changes to cellular constituents (Walters *et al.*, 2010).

Structural changes associated with oxidation are reduced membrane fluidity, altered folding of DNA, lost elasticity of proteins and increased brittleness of the cellular matrix. Molecules oxidation leads to either smaller molecules with reactive carbonyl or nitrogen groups that easily diffuse through cells, or adducts between carbohydrates, proteins and nucleic acids that cause intermolecular cross-linking and further degrade into advanced glycation end-products (Walters *et al.*, 2010).

### **Biochemical Manifestation of Seed Deterioration**

Seed deterioration is associated with various cellular, metabolic and chemical alterations including chromosome aberrations and damage to the DNA, impairment of RNA and protein synthesis, changes in the enzymes and food reserves and loss of membrane integrity (Kibinza *et al.*, 2006). Some of the major physiological and biochemical events of deterioration are presented below.

#### **Membrane Degradation**

It is extensively consented that loss in cellular membrane integrity is one of the primary causes for loss of viability. Under harsh storage conditions loss in membrane permeability leads to increased leaching of seed constituents and hence loss in

viability. During seed deterioration, membrane degradation increase electrolyte leakage. Decline in seed germination, field emergence and seedling vigor is associated with high level of electrolytes leakage.

Alterations of membrane systems, such as the tonoplast, plasmalemma and endoplasmic reticulum, result in diminishing of normal cell function and energy production. Membrane deterioration and loss of permeability occur at an early stage during the seed deterioration.

#### **Enzymes Alterations**

Enzyme alterations, such as reduced activity of lipase, ribonuclease, acid phosphatase, protease, diastase, catalase, peroxidase,  $\alpha$  and  $\beta$  amylase, DNase and dehydrogenase enzymes. ROS and hydrogen peroxides are produced from several metabolic reactions and could be destroyed by the activity of scavenger enzymes like catalase and peroxides. Peroxides activity decreases substantially with ageing. Due to this seeds become more sensitive to the effects of oxygen and free radicals in membrane unsaturated fatty acids and produce lipid peroxidation products such as monaldehyde and lipid conjugants.

#### **Changes in Cell Chemical Constituents**

In deteriorate seeds significant decrease in protein, oil content and total sugars and increase in free fatty acids and reducing sugars has been studied. Verma *et al.* (2003) showed that carbohydrates increased with decrease in protein content in deteriorated seeds. Some studies indicated that oligosaccharide which has been associated in stabilizing membranes decreased during storage.

**Reduced Metabolic Activity**

High relative humidity hastens deterioration and results in reduction of nucleic acids with increased storage period. Metabolic activities of seeds were low in non-viable seeds than in viable seeds. Long term storage decreases the ability to form nucleic acids and nucleotides.

**Free Radical Damage**

Deterioration is partially associated to the accumulation of free-radicals produced by the metabolic process. Seed storage subjects lipids to slow consistent attack by oxygen, forming hydrogen peroxides, other oxygenated fatty acids and free radicals. The free radicals are unstable and may react and damage nearby molecules. Oxygenated fatty acids in the absence of enzymes activity in the dry seed would accumulate and damage cellular components and leads to deterioration of seeds. Lipid peroxidation and free radicals formation are the major causes for the deterioration of oil seeds in storage.

**Chromosome Aberrations**

One of the changes linked with seed ageing is aberration of chromosomes, sometimes pertained to as mutagenic effects. Some of the chromosome alterations in seeds comprise fragmentation, bridges, fusion, ring formation of chromosomes and variations in nuclear size.

Some other causes of deterioration are:

- Degradation of functional structures,
- Biochemical changes resulting in lower levels of ATP,
- Decline in sugar content,
- Inability of ribosomes to dissociate,

- Enzyme degradation and inactivation (amylase, dehydrogenase, oxidases, phospholipase, glutamic acid decarboxylase),
- Formation and activation of hydrolytic enzymes,
- Starvation of meristematic cells,
- Increases in seed leachates and free fatty acid content,
- Reduced respiration, and
- Accumulation of toxic compounds.

Moreover the main cause of seed damage, lipid peroxidation causes initial biochemical changes in seed that can be observed during storage. Autooxidation of lipids and increase in the content of free fatty acids throughout storage period are the main reasons for rapid deterioration of seed of oil plants (Balesevic-Tubic *et al.*, 2005).

In sunflower seeds, loss of viability is associated with an accumulation of malondialdehyde (MDA), suggesting that seed deterioration is accompanying with lipid peroxidation related to a decline in the efficiency of the antioxidant defense system (Kibinza *et al.*, 2006).

Membrane disruption is one of the primary reasons attributed to seed deterioration. As a result, seed cells are not capable to hold their normal physical condition and function. Causes of membrane disruption are enhancing in free fatty acid level and free radicals productivity by lipid peroxidation (Ghassemi-Golezani *et al.*, 2010). Lipid peroxidation can result in not only destruction of the lipid itself, but also damage to cell membranes and other cellular components. Free fatty acid can damage lipid bilayer

particularly of mitochondria leading to reduce energy production and free radicals have potential to damage membrane, DNA, enzymes, protein and ultimately cellular repair mechanism (Ghassemi-Golezani *et al.*, 2010).

### **Factor Affecting Deterioration**

The rate of seed deterioration is highly influenced by environmental (temperature, relative humidity and seed moisture content) and biological factors (such as fungi that create their own biological niche) (Ghassemi-Golezani *et al.*, 2010). Seed longevity is determined by seed moisture, temperature and seed attributes that are influenced by genetic and environmental interactions during seed maturation, harvesting and storage (Walters *et al.*, 2010).

Several other factors such as environmental conditions during seed producing stage, pests, diseases, seed oil content, storage longevity, mechanical damages of seed in processing, fluctuations in moisture (including drought), weathering, nutrient deficiencies, packaging, pesticides, improper handling, drying and biochemical injury of seed tissue can affect vigor of seeds (Krishnan *et al.*, 2003; Marshal and Levis, 2004; Astegar *et al.*, 2011; Simic *et al.*, 2007).

### **Kind/variety of the Seed**

The seed storability is considerably determined by the kind or variety of seeds. Some seeds are naturally short-lived, e.g., onion, soybeans, peanuts, etc., whereas some seeds like, tall fescue and annual rye grass, appear very similar but differ in storability. Genetic make-up of varieties also influences storability.

### **Genotypic Factors**

Some types of seeds are inherently long lived; others are short lived, while others have an intermediate life span owing to their differences on genetic makeup.

### **Initial Seed Quality**

High initial viability of seeds maintains their quality in storage longer than those with less initial viability. Vigorous and undeteriorated seeds can store longer than deteriorated seeds. Seeds that have been broken, cracked, or bruised due to handling deteriorate more rapidly in storage than undamaged seeds. Cracks in seeds serve as entrance to pathogens causing consequent deterioration. Seeds that have been developed under environmental stress conditions (such as drought, nutrient deficiency and high temperatures) become more susceptible to rapid deterioration.

### **Effect of Temperature**

High temperature hastened the rate of these biochemical processes triggering more rapid deterioration that resulted in rapid losses in seed having high moisture content (Shelar *et al.*, 2008). Seeds sensitivity to high temperatures is strongly dependent on their water content, loss of viability being quicker with increasing moisture content (Kibinza *et al.*, 2006). Temperature is important because it influences the amount of moisture and also enhances the rate of deteriorative reactions occurring in seeds as temperature increases.

### **Effect of Moisture Content**

Deteriorative reactions occur more readily in seeds at higher moisture content and subsequently, this condition constitute hazard to the longevity of seed survival (Vashisth and Nagarajan, 2009).

Seeds stored at high moisture content demonstrate increased respiration, heating, and fungal invasion resulting in reduced seed vigor and viability. After physiological maturity the rate of seed quality loss depends on the degree of unfavorable environmental conditions surrounding the seed. Environmental moisture, predominantly intermittent or prolonged rainfall, during the post-maturation and pre-harvest period, is quite detrimental to seed quality and cause rapid deterioration. When exposed to humid conditions (heavy rain), dried seeds can absorb enough moisture to reach 27% and subsequently expand in volume. At this moisture level, seed respiration is hastened. Cotyledonary reserves will be consumed, not only by the seed itself, but also by fungi allied with the seed. It has been reported that seed moisture content of about 6-8% is optimum for maximum longevity of most crop species. Below 4-6% seed moisture content lipid autoxidation becomes a damaging factor and seeds become more susceptible to mechanical damage.

The moisture content of seed during storage is the most persuasive factor affecting the longevity.

Storing seeds at high moisture content enhances the risk of quicker deterioration at shorter time. Seeds are hygroscopic in nature; they can pick up and releases moisture from and to the surrounding air. They absorb or lose moisture till the vapor pressure of seed moisture and atmospheric moisture reach equilibrium (Shelar *et al.*, 2008). Control of relative humidity is the most important because it directly influences the moisture content of seeds in storage as they come to equilibrium with the amount of moisture surrounding them; a concept

known as equilibrium moisture content. The lower the moisture content, the longer seeds can be stored provided that the moisture level can be controlled all through the storage period.

#### **Effect of Organisms Associated with Seeds**

Organisms associated with seeds in storage are bacteria, fungi, mites, insects and rodents. The activity of these entire organisms can lead to damage resulting in loss of vigor and viability or, complete loss of seed.

**Bacteria and fungi:** There are several factors which favor infection fungi and promote their infestation such as moisture content of seed and interspace relative humidity, temperature, pre-storage infection and storage pest. Most storage fungi belong to *Penicillium* and *Aspergillus* genera. They induce seed deterioration by producing toxic substances that destroy the cells of seeds. Mechanically damaged seed allow quick and easy access for mycroflora to enter the seed (Shelar *et al.*, 2008). To minimize the risk of fungi invasion, seeds have to be stored at low moisture content, low temperature, and RH.

Researches show that all storage fungi are completely inactive below 62% relative humidity and show very little activity below about 75% relative humidity upwards, the amount of fungi in a seed often shows an exponential relationship with relative humidity. The storage bacteria require at least 90% relative humidity for growth and therefore only become significant under conditions in which fungi are already very active.

**Insect and Mites:** There is no insect activity at seed moisture contents below 8%, but if grain is infected, increased activity may generally be expected up to about 15% moisture content. The optimum temperature for insect activity of storage



insects ranges from 28 to 38°C. The temperatures below 17 to 22°C are considered unsafe for insect activity. Although it is usually preferable to control insect and mite activity by the manipulation of the seed environment, i.e., use of fumigants and insecticides. The main problem of chemical control is the adverse effect of chemicals on seed viability and vigor, and some of them are dangerous to handle. However, fumigants which have been used successfully include methyl bromide, hydrogen cyanide, phosphine, ethylene dichloride and carbon tetrachloride in 3:1 mixture, carbon disulphide and naphthalene. Insecticides – used in seed storage include DDT, lindane and Malathion.

#### **Provenance**

Seeds obtained from different sources may show differences in viability and storability. Nevertheless, the seed begins its existence before it harvest and it is expected that seeds harvested in different pre-harvest condition.

#### **Fluctuating Environmental Conditions**

Fluctuating environmental conditions are harmful for seed viability. Rapid changes in seed moisture content and temperature cause deleterious effect.

#### **Oxygen Pressure**

Recent researches on the role of a gaseous environment on seed viability indicate that increases in pressure of oxygen incline to decreases the viability period.

#### **Other Factors**

Factors besides those discussed above that affect seed storage life are the direct sunlight on the seed, number of times and kind of fumigation, effect of seed treatment, etc.

#### **Symptoms of Seed Deterioration**

Seed deterioration is an inexorable, irreversible

degenerative change in the quality of a seed. It is observable in their dropped performance during germination such as delayed seedling emergence, slower rate of seedling growth and development, loss of the capacity to germinate was the final phase of deterioration process and final indication of vigor loss. Deteriorated seeds are also diminished resistance to environmental stresses during germination and early seedling growth.

Some of the symptoms of the deteriorated seed are:

#### **Morphological Changes**

Changes in seed coat color are apparently owing to oxidative reactions in the seed coat that are enhanced under conditions of high temperature and relative humidity.

#### **Ultra Structural Changes**

By electron microscopy examination, two broad patterns of coalescence of lipid bodies and plasmalemma extraction from the cell wall related with deterioration have been observed. Both of these events influence cell membrane integrity.

#### **Cell Membrane Changes**

As seeds deteriorate, their incapability to retain cellular constituents decreased which was attributed to cell membrane disruptions associated with the loss of membrane phospholipids. Some consequence of membrane damage includes:

- Breaks in the plasmalemma structure and its contraction from the cell wall,
- Fragmented endoplasmic reticulum lacking of polyribosomes,
- Lack of dicytosomes,

- Monosomes arbitrarily dispersed in the cytoplasm,
- Disintegration of mitochondria and plastids,
- Coalescence of lipid droplets,
- Condensation of chromatin and lobed nucleus, and
- Lyses of membranes of lysomic structures.

### **Loss of Enzyme Activity**

The most sensitive test for measuring early seed deterioration is tetrazolium (TZ) and glutamic acid decarboxylase activity test. Other oxidases enzymes such as catalase, peroxidase, amylase and cytochrome oxidase also correlated with seed deterioration.

### **Reduced Respiration**

As seeds deteriorate, respiration becomes gradually pathetic and eventually leads to loss of germination. Reduction in respiration rate is closely associated with seed deterioration.

### **Increase in Seed Leachates**

Increase in leachate content when soaked in water is often noticed symptom of deteriorated seeds. The leachate concentration can be measured by electrical conductance methods and by determines the soluble sugar content of the leachate.

### **Increase in Free Fatty Acid Content**

The hydrolysis of phospholipids leads to the release of glycerol and fatty acids, and this reaction hastens with increasing seed moisture content. The frequent accumulation of free fatty acids concludes in a decline in cellular pH and is detrimental to normal cellular metabolism.

## **Methods for Testing Seed Deterioration**

### **Germination Test**

It is an analytical procedure to evaluate seed germination under standardized, favorable conditions. Standard germination testing includes media, temperature, moisture, light, dormancy breaking and germination counting standard for various crop seeds.

### **Tetrazolium (TZ) Test**

TZ test is extensively accepted as an accurate mean of estimating seed viability. This method was developed by Professor Georg Lakon in the early 1940s. It is quick method to estimate seed viability (Copeland and McDonald, 2001). This test distinguishes between viable and dead tissues of the embryo on the basis of their relative respiration rate in the hydrated state.

### **Electrical Conductivity Test**

As seed deterioration progresses, the cell membranes become less rigid and become more water permeable. It allows the cell contents to leakage into solution with the water and increasing electrical conductivity. It provides a rapid indication of seed viability for seed lots.

### **Vital Coloring Test**

The principle of this method is the differential coloration of live against dead tissues when exhibited to certain dyes such as sulfuric acid, indigo carmine and aniline dyes. These dyes stain the dead tissue blue and the live tissue leftovers unstained. This method is particularly useful for determining viability of tree seeds.

### **Enzyme Activity Test**

These methods measure enzyme activity (such as lipase, amylase, diastase, catalase, peroxidase and dehydrogenase) of imbibed seeds as an indication of their viability.

**Other Tests**

Other testing methods are free fatty acid test, hydrogen peroxide test, indoxyl acetate test, fast green test, ferric chloride test, sodium hypochlorite test, excised embryo test and X-ray test. These methods were discussed by Copeland and McDonald (2001).

**Detrimental Effects of Seed Deterioration**

Some probable consequences of deteriorative changes in seeds are:

- Decreased percent germination;
- Reduction in vigor and viability;
- Degradation of cellular membranes and loss of permeability control;
- Increased solute leakage;
- Impairment of energy-yielding and biosynthetic mechanisms;
- Reduced biosynthesis and respiration;
- Reduced germination rate and early seedling growth;
- Reduced rate of plant growth and development;
- Reduced storage potential;
- Decreased growth uniformity;
- Increased susceptibility to environmental stresses, especially during germination, emergence, and early seedling development;
- Reduced tolerance under adverse conditions
- Decreased yield;
- Decreased emergence percentage;
- Increased percentage of abnormal seedling;
- Loss of the capacity to germinate; and

- Loss in seed weight (Mohammadi *et al.*, 2011; Ghassemi-Golezani *et al.*, 2010; Astegar *et al.*, 2011; Farhadi *et al.*, 2012; Biabani *et al.*, 2011; Shelar *et al.*, 2008).

It also results in decreasing the quality of seed produced, such as purple stained seeds (contaminated by fungi), wrinkled seeds, fissures in the seed coat, insect damaged seeds, discoloration of seed, etc.

**CONCLUSION**


Seed quality, germination, vigor and viability are highly influenced by environmental factors in field and storage. There was a distinct reduction in yield, seedling growth, loss of capacity to germinate and increased susceptibility to environmental stresses which cause numerous harmful effect on seed quality.

**REFERENCES**

1. Astegar Z R, Sedghi M and Khomari S (2011), "Effects of Accelerated Aging on Soybean Seed Germination Indexes at Laboratory Conditions", *Not Sci Biol.*, Vol. 3, No. 3, pp. 126-129.
2. Balesevic-Tubic S, Malenèiæ D, Tatiæ M and Miladinovic J (2005), "Influence of aging process on biochemical changes in sunflower seed", *HELIA*, Vol. 28, No. 42, pp. 107-114.
3. Bhatia V S, Yadav S, Jumrani K and Guruprasad K N (2010), "Field deterioration of Soybean seed: Role of oxidative stress and antioxidant defense mechanism", *Jour. Plant Biol.*, Vol. 32, No. 2, pp. 179-190.
4. Biabani A, Boggs LC, Katozi M and Sabouri H (2011), "Effects of seed deterioration and inoculation with *Mesorhizobium cicerion*

- yield and plant performance of chickpea”, *Australian Jour. of Crop Science*, Vol. 5, No. 1, pp. 66-70.
5. Copeland L O and McDonald M B (2001), *Principles of Seed Science and Technology*, Kluwer academic publishers, Boston, p. 330.
  6. Farhadi R, Rahmani M R, Salehi balashahri M and Sadeghi M (2012), “The effect of Artificial ageing on germination components and seedling growth of Basil (*Ocimum basilicm* L.) seeds”, *Journal of Agriculture and Food Technology*, Vol. 2, No. 4, pp. 69-72.
  7. França-Neto J B, Pádua G P, Carvalho M L M D E, Costa O, Brumatti P S R, Krzyzanowski F C, Costa N P da, Henning A A and Sanches D P (2005), “Semente esverdeada de soja e sua qualidade fisiológica” Londrina, *Embrapa Soja*, p. 8, *Embrapa Soja. Circular Técnica*, 38.
  8. Ghassemi-Golezani K, Bakhshy J, Raey Y and Hossainzadeh-Mahootchy A (2010), “Seed Vigor and Field Performance of Winter Oilseed Rape (*Brassica napus* L.)”, *Cultivars. Not. Bot. Hort. Agrobot. Cluj.*, Vol. 38, No. 3, pp. 146-150.
  9. Jatoi S A, Afzal M, Nasim S and Anwar R (2001), “Seed deterioration study in pea, using accelerated ageing techniques”, *Pakistan Jour. of Biological Sciences*, Vol. 4, No. 12, pp. 1490-1494.
  10. Kapoor N, Arya A, Siddiqui M A, Kumar H and Amir A (2011), “Physiology and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.)”, *America Jour. of Plant Physiology*, Vol. 6, No. 1, pp. 28-35.
  11. Kapoor R, Arya A, Siddiqui, MA, Amir A and Kumar H (2010), “Seed Deterioration in Chickpea (*Cicer arietinum* L.) under Accelerated Ageing”, *Asian J Plant Sci.*, Vol. 9, No. 3, pp. 158-162.
  12. Khatun A, Kabir G, Bhuiyan MAH (2009), “Effect of harvesting stages on the seed quality of lentil (*Lens culinaris* L.) during storage”, *Bangladesh Jour. Agril. Res.*, Vol. 34, No. 4, pp. 565-576.
  13. Kibinza S, Vinel D, Côme D, Bailly C and Corbineau F (2006), “Sunflower seed deterioration as related to moisture content during ageing, energy metabolism and active oxygen species scavenging”, *Physiologia Plantarum*, Vol. 128, No. 3, pp. 496-506.
  14. Krishnan P, Nagarajan S, Dadlani M and Moharir A V (2003), “Characterization of wheat (*Triticum aestivum* L.) and soybean (*Glycine max* L.) seed under accelerated aging condition by proton nuclear magnetic spectroscopy”, *J Seed Sci Technol.*, Vol. 31, pp. 541-550.
  15. Mahesha C R, Channaveeraswami A S, Kurdikeri M B, Shekhargouda M and Merwade M N (2001a), “Seed maturation studies in sunflower genotypes”, *Seed Res.*, Vol. 29, No. 1, pp. 95-97.
  16. Marshal A H and Levis D N (2004), “Influence of seed storage conditions on seedling emergence, seedling growth and dry matter production of temperate forage grasses”, *J Seed Sci Technol.*, Vol. 32, pp. 493-501.

17. Mohammadi H, Soltani A, Sadeghipour H R and Zeinali E (2011), "Effect of seed aging on subsequent seed reserve utilization and seedling growth in soybean", *Inter. Jour. of Plant Production*, Vol. 5, No. 1, pp. 65-70.
18. Mosavi Nik S M, Gholami tilebeni H, Kord firouz jae GH, Sadeghi M and Sedighi E (2011), "Free fatty acid and electrical conductivity changes in cotton seed (*Gossypium hirsutum*) under seed deteriorating conditions", *International Journal of Agri Science*, Vol. 1, No. 2, pp. 62-66.
19. Pádua GP, França-neto JB, Carvalho M L M, Krzyzanowski F C and Guimarães R M (2009), "Incidence of green soybean seeds as a function of environmental stresses during seed maturation", *Revista Brasileira de Sementes*, Vol. 31, No. 3, pp. 150-159.
20. Shelar V R (2008), "Role of mechanical damage in deterioration of soybean seed quality during storage- a review. *Agric. Rev.*", Vol. 29, No. 3, pp. p177-184.
21. Shelar V R, Shaikh R S and Nikam A S (2008), "Soybean seed quality during storage: A review", *Agric. Rev.*, Vol. 29, No. 2, pp. 125-131.
22. Simic B, Popoviae R, Sudaric A, Rozman V, Kalinovic I and Cosic J (2007), "Influence of Storage Condition on Seed Oil Content of Maize, Soybean and Sunflower", *Agriculturae Conspectus Scientificus*, Vol. 72, No. 3, pp. 211-213.
23. Tilebeni G H and Golpayegani A (2011), "Effect of seed ageing on physiological and biochemical changes in rice seed (*Oryza sativa* L.)", *International Journal of AgriScience*, Vol. 1, No. 3, pp. 138-143.
24. Vashisth A and Nagarajan S (2009), "Germination Characteristics of Seeds of Maize (*Zea mays* L.) Exposed to Magnetic Fields under Accelerated Ageing Condition", *Journal of Agricultural Physics*, Vol. 9, pp. 50-58.
25. Verma S S, Tomer R P S and Verma U (2003), "Loss of viability and vigor in Indian mustard seeds stored under Ambient conditions", *Seed Res.*, Vol. 31, No. 1, pp. 98-101.
26. Walters C, Ballesteros D and Vertucci V A (2010), "Structural mechanics of seed deterioration: Standing the test of time", *Plant Science*, Vol. 179, pp. 565-573.



**International Journal of Life Sciences Biotechnology and Pharma Research**

**Hyderabad, INDIA. Ph: +91-09441351700, 09059645577**

**E-mail: editorijlbpr@gmail.com or editor@ijlbpr.com**

**Website: www.ijlbpr.com**

